

# CHAMPION- AF

Clinical trial to determine if left atrial appendage closure with the WATCHMAN FLX device is a reasonable alternative to non-vitamin K oral anticoagulants (NOACs) in patients with non-valvular Atrial Fibrillation

Primary Inclusion	Primary Exclusion
<input type="checkbox"/> Age > 18 years old	<input type="checkbox"/> Requires long-term anticoagulation therapy for reasons other than AF-related stroke risk reduction
<input type="checkbox"/> Documented non-valvular atrial fibrillation (i.e. atrial fibrillation, in the absence of moderate or greater mitral stenosis or a mechanical heart valve)	<input type="checkbox"/> The subject had or is planning to have any cardiac or non cardiac intervention or surgical procedure within 30 days prior to or 60 days after implant (including, but not limited to: cardioversion, percutaneous coronary intervention, cardiac ablation, cataract surgery, etc.)
<input type="checkbox"/> The subject has a calculated CHA2DS2-VASc score of 2 or greater for men and 3 or greater for women	<input type="checkbox"/> Indicated for chronic P2Y12 platelet inhibitor therapy
<input type="checkbox"/> Suitable for the protocol defined pharmacologic regimens in both the test and control arms	<input type="checkbox"/> History of atrial septal repair or has an ASD/PFO device <input type="checkbox"/> Subject has an implanted mechanical valve prosthesis in any position
	<input type="checkbox"/> LVEF < 30%

*PI: Albert Lin, MD*

*Study Coordinator: Amy Carswell*

*Pager: 5-7501, Phone: 6-7554, amy.carswell@nm.org*