

COVID-19 Inpatient Orders and VTE Orders

Impacted Locations	All
Intended Audience	Inpatient Physician and APP

COVID-19 Inpatient Orders and Admission VTE Orders

Scenario

When merging order sets, such as **Admission** and **COVID-19 Inpatient Orders**, both containing VTE, the admission VTE section will be required, as the VTE Order must be signed before it addresses the requirement.

Handling this scenario:

- **Option 1:** Complete the COVID-19 Inpatient Orders, addressing VTE Orders prior to opening the Admission Order Set.
- **Option 2:** From the Admission Order Set, select **Chemical Prophylaxis Contraindicated**, and choose the reason of **Other (see comments)**, COVID-19.

Option 2

From the Prophylaxis, Chemical VTE Prophylaxis section:

1. Select **Chemical Prophylaxis Contraindicated**.

▼ Prophylaxis

▼ Chemical VTE Prophylaxis

Patient currently has an active chemical prophylaxis order

- CDH/DELNOR - (VTE) Prophylaxis Guidelines: Adult
- KH - (VTE) Prophylaxis Guidelines: Adult
- VWH - (VTE) Prophylaxis Guidelines: Adult
- NMH/LFH - (VTE) Prophylaxis Guidelines: Adult

heparin bridge to enoxaparin (LOVENOX) - when scheduled dose of enoxaparin is GREATER than 16 hours from time of order

enoxaparin (LOVENOX) injection
40 mg, Subcutaneous, DAILY

heparin (porcine) injection
5,000 Units, Subcutaneous, 3 TIMES DAILY

fondaparinux (ARIXTRA) - For History of HIT or Heparin Allergy

Chemical Prophylaxis Contraindicated
⚠ Routine, UNTIL DISCONTINUED, starting today at 1559, Until Specified

2. From the order composer, select the reason **Other (see comments)**

Reason for no chemoprophylaxis:

Coagulopathy (INR>2 or PT>18) Likely Procedure in 12 Hours

Risk of Active Hemorrhage Post-Operative Bleeding Concerns

Risk of Intracranial Hemorrhage Hemophilia Platelets (<50,000)

Platelet Dysfunction Hypothermia (T < 95 F) Hypertension (SBP > 200)

Liver Dysfunction (MELD > 10) Postpartum Low Risk of VTE Low Risk of VTE

Therapeutic Anticoagulation Admission to OB **Other (see comments)**

Other comments: COVID-19

3. In **Other comments**, enter **COVID-19**.
4. From the **Medications, VTE Prophylaxis** section, select the VTE orders.